

METHOD FOR EVALUATING AND AFFECTING MALE FERTILITY

FIELD OF THE INVENTION

[0001.] The present invention relates to a sperm protein which can be used for evaluating, inhibiting, and/or enhancing male fertility, as well as antibodies to the sperm protein.

INS
C1
The present application is a continuation in part of application Serial No. 09/123,492, filed July 28, 1998, Serial No. and PCTUS9701725 filed January 29, 1998, which is a continuation in part of application Serial No. 08/592,677, filed January 29, 1996 and now abandoned, both of which are hereby incorporated by reference in the entirety. The present application is based on and claims priority from provisional application No. 60/082,753, filed April 23, 1998.

BACKGROUND OF THE INVENTION

[0002.] Sperm production in the testis of human males is far less efficient than sperm production in other mammals, such as rat, rabbit and monkey (Amann, 1970) due to an increased rate of germ cell atresia. Together with this is the fact that a high incidence of sperm in the ejaculate of a fertile man is morphologically abnormal (Wyrobek et al., 1982). Thus, there is a heightened awareness of the possibility that the quantity and quality of sperm in the ejaculates of men are declining because of environmental influences (Sharpe, 1993). A toxicant-induced alteration in the process of sperm maturation during sperm transit through the epididymis, the organ in which sperm acquire fertilizing ability, could render a man infertile. It has been hypothesized that specific proteins

are added to sperm in the epididymis which confer fertility. Recently, Klinefelter et al., in Journal of Andrology 15(4), 318-327 (1994) demonstrated that an 18 kD epididymal sperm surface protein, presumably a plasma membrane protein, was well correlated with fertility, although it was not believed that this protein was predictive of fertility.

[0003.] There continues to be great interest in developing new and improved contraceptives. New contraceptives should be superior to existing products, e.g., oral contraceptives used by millions of women over the last 30 years are not only safe and effective but even protect women against some cancers. However, other methods of contraception are still needed by many segments of the world's population, as many women do not have reliable access to oral contraceptives, or may suffer adverse reactions to the hormones use in oral contraceptives.

[0004.] Additionally, fertility testing is becoming more widespread as increasing numbers of apparently infertile couples seek medical assistance in conception. Because reproductive abnormalities of both sexes may affect fertility, assessing male fertility is common in fertility evaluations. While the most common starting point for evaluation of male fertility is an assessment of the sperm count in semen, also important to fertility is sperm motility. Therefore, in male fertility analyses, sperm motility has also been a factor.

[0005.] Currently available techniques for measuring sperm count and sperm motility are microscopic in nature. A

[0006.] Attempts to develop biochemical assays of semen have not resulted in simple procedures which may be performed in either the physician's office or a dedicated semen evaluation lab. Most biochemical markers have failed to demonstrate correlations with sperm number, motility, or fertility. Activity of fumarase, an enzyme present in semen, has been found to correlate to both sperm count and percentage motility, Crabbe, J. Reprod. Fert. 51:73-76 (1977). Crabbe measured fumarase activity by spectrophotometric measurements. Unfortunately, spectrophotometric assays are not generally suitable for office assays because of the cost of these specialized devices as well as the training required for accurate and reproducible operations.

[0007.] Dorian, in U.S. Patent No. 5,434,057, expanded on Crabbe's method by providing devices for assessing sperm number and motility in semen samples comprising a solid support having a carrier matrix containing a fumarase substrate and malate dehydrogenase. The sample is applied to the carrier matrix and a visual signal is detected from the

solid support resulting from metabolism of the fumarase substrate by fumarase in the sample. While this assay detects motile sperm in a semen sample, there is not method for inhibiting fertility nor of selecting out the most fertile sperm in a sample.

[0008.] Feuchter et al., in U.S. Patent No. 5,250,417, disclose a method for detecting the ability of sperm to undergo the acrosome reaction to permit determination of the fertility of male mammals. The acrosome reaction is a process by which sperm release hydrolytic enzymes that degrade the zona pellucida, which must be penetrated to enable the sperm and ovum to come into contact, fuse, and complete the fertilization process.

[0009.] In recent years, other studies have targeted different proteins associated with sperm in an attempt to provide new contraceptive alternatives. Major research efforts involve immunological approaches to fertility control. The development of contraceptive vaccines is directed towards the immunoneutralization of reproductive processes or interfering with fertilization by inducing antibodies against oocytes and spermatozoa. Several sperm antigens shown to have high immunocontraceptive potential are human sperm membrane antigen (SP-10) and guinea pig sperm membrane protein (PH-20). SP-10 is a sperm membrane specific antigen of 24-24 kD which was isolated using a monoclonal antibody (MHS-10) that cross-reacts with the entire acrosomal region. It is associated

09554-000

with the outer aspect of the inner acrosomal membrane and the inner aspect of the outer acrosomal membrane of mature human sperm. It has been reproduced recombinantly in an Escherichia coli expression system.

[0010.] PH-20, a guinea pig sperm protein of 64 kD, is present on both the plasma membrane and inner acrosomal membrane of sperm. It is essential for adhesion of sperm to the zona pellucida, the initial step in the fertilization process. Active immunization with PH-20 causes infertility in both male and female guinea pigs for a period ranging from six to fifteen months.

[0011.] O'Rand et al., in U.S. Patent No. 5,175,148, disclose a sperm antigen corresponding to a sperm autoantigenic epitope which can be used as immunocontraceptive agent as well as for diagnosing autoimmune infertility. The synthetic peptide corresponds to an autoantigenic epitope of rabbit sperm membrane autoantigen.

[0012.] Several other antigens with good immunocontraceptive potential have been identified and investigated in laboratory animals, including lactic dehydrogenase-x, an isoenzyme of lactic dehydrogenase confined to male germ cells. A synthetic peptide based upon a portion of this antigen has been shown to reduce fertility in laboratory animals. Unfortunately, most studies have found that the rate and duration of the immunocontraceptive effects are less than acceptable. A problem in immunological approaches to antifertility research

0973254-010301

is the need for a safe, effective adjuvant and suitable animal models for evaluating the efficacy and safety of methods.

[0013.] Although most contraceptive research has been directed to use in females, there is an interest in male fertility control both from a scientific as well as a biological viewpoint. Many compounds have been identified as having male antifertility activity in various species, e.g., gossypol, 5-thio- D-glucose, and 6-chlorodeoxyglucose. Studies have also been conducted on the use of androgens to control male fertility. Unfortunately, most compounds identified as useful in controlling male fertility appear either to have irreversible antifertility effects, to be inherently toxic, or to affect libido. It has been demonstrated that sperm count could be depressed in men injected with large doses of androgens. However, there are still questions about the potential utility of androgens as male antifertility agents. The ideal male contraceptive would produce azoospermia without compromising libido or sexual potency, and would be reversible.

[0014.] While numerous sperm proteins (Primakoff et al., 1997; Burks et al., 1995; Wei et al., 1994; o'Rand et al., 1984; Lea et al., 1996; O'Rand et al. 1996; Amman et al., 1998a,b; Hammerstedt et al., 1997; Cohen et al., 1996) as well as seminal plasma proteins (Killian et al., 1996; Peknicova et al., 1997) have been associated with fertility over the years, in most cases they have been identified based on the

097534-010201

demonstration of sperm-egg binding in vitro, a system that differs considerably from the natural environment of sperm.

[0015.] In recent years several proteins have been linked in some way or another to fertilizing capacity of sperm. While most of these proteins are associated with the sperm membrane, some have been identified in seminal plasma. Two of the sperm proteins are quite large. The first of these is a 64 kD sperm membrane protein (PH-20) which was found to localize over the arosome of guinea pig sperm and believed to function during fertilization (Hunnicuttt et al., 1996). Antibody to PH-20 has been shown to prevent binding of sperm to the zona pellucida, and males were rendered infertile when PH-20 was administered as an immune vaccine (Primakoff et al., 1997). However, PH-20 has only been described in the guinea pig, and the PH-20 vaccine severely compromises spermatogenesis in older animals, rendering the effect irreversible. In addition, a 95 kD mouse sperm protein was identified as a phosphotyrosine protein ligand for ZP3, a glycoprotein in the extracellular matrix of the egg (Burks et al., 1995). Specific peptide fragments of this protein blocked binding of sperm to the zone pellucida in vitro, but inhibition of fertility in vivo has not been demonstrated.

[0016.] Several small sperm proteins have been linked to fertility. Rabbit sperm autoantigen I (RSAI), now referred to as SP-17 (Lea et al., 1996), is a 24 kD protein unique to the testis. This protein appears to be common to rabbit, mouse,

09752344-040304

and humans, as evidenced by cDNA sequence homology. Monospecific antibodies to this protein have not been used to demonstrate antifertility effects *in vivo*. There is no protein homology based on cDNA and deduced amino acid sequence between SP-17 and SP22.

[0017.] Protein D and E are secreted by the epithelium of the epididymis. These proteins have been linked to fertility as the plasma membrane of the oocyte exposes protein D/E binding domains during sperm-egg fusion (Cohen et al., 1996). Antibody directed against the D/E complex significantly inhibited sperm penetration of zona-free eggs *in vitro* (Cuasnicu et al., 1990), but no fertility role has yet been demonstrated *in vivo*. While these proteins have molecular weights of 26 and 32 kD, respectively, the sequence of SP22, the protein of the present invention, is not related to that of proteins D or E.

[0018.] Wei et al., 1994, recovered a 17 kD protein from a detergent extract of human sperm. Antibody to this glycoprotein localized over the head of sperm multiple species, but staining was localized over the entire head of the sperm as well as over the principal piece of the tail of the sperm, suggesting a lack of specificity. Thus, while the antibody inhibited fertility *in vivo*, much of this inhibition appears to result from nonspecific binding. Unfortunately, adequate control data were not shown. The 17 kD protein described by Wei et al. (1994) was never identified in the

0975344-010304

profile of proteins in the original detergent extract. Thus, it is impossible to determine the relationship of this protein relative to others reported in the literature. It seems that this may be similar to the more recently-described SP-17 described by Lea et al., 1996, but this is impossible to determine, as Wei et al. did not provide amino acid sequence data.

[0019.] O'Rand et al. described DNA encoding a mammalian Sp17 protein or antigenic peptides which are fragments thereof. These proteins and fragments are said to be useful as immunocontraceptive agents and for diagnosing autoimmune fertility.

[0020.] Two proteins identified in the seminal fluid, the sperm-free fraction of semen, have been linked to fertility. The first is a 17 kD protein referred to as ACR.3 (Capkova and Peknicova, 1997). This protein is a coating protein rather than an integral membrane protein and is apparently involved in mediating sperm binding to the zona pellucida, as addition of purified ACR.3 to normal sperm actually diminishes their capacity to bind to the zone. The function of these seminal plasma proteins may be to prevent premature capacitation and binding of sperm to the egg. Such membrane stabilizers may be critical to normal fertilization *in vivo*, but are not diagnostic of fertilizing potential. In fact, antibody to ACR.3 does not inhibit fertilization. The second seminal plasma protein that has been associated with the fertility of

00000449260

bull sperm is a 265 kD protein (Killian et al., 1996), now known as prostaglandin D-synthase (Gerena et al., 1998). This protein may have more direct influence on fertility, as the addition of seminal plasma from higher fertility bulls increased the ability of sperm from lower fertility bulls to penetrate the egg *in vitro*. However, neither the direct addition of purified 26 kD protein, nor *in vivo* function tests, have been performed.

SUMMARY OF THE INVENTION

[0021.] It is an object of the present invention to overcome the aforesaid deficiencies in the prior art.

[0022.] It is another objection of the present invention to provide a means for predicting male fertility in animals as well as in human males.

[0023.] It is another object of the present invention to screen for environmental endocrine disruptors, as endocrine disruption can lead to male-mediated infertility.

[0024.] It is yet another object of the present invention to screen animals and humans exposed to known or suspected endocrine disruptors for fertility.

[0025.] It is still another object of the present invention to select sires for artificial insemination who are good candidates for providing sperm for insemination.

[0026.] It is a further object of the present invention to screen human semen for fertility prior to undertaking assisted reproductive technology techniques to improve the success of

092524-010304

these techniques.

[0027.] It is a further object of the present invention to improve fertility in males who fail to express a sufficient amount of SP22, formerly known as SP-16, in sperm.

[0028.] It is another object of the present invention to provide a reversible male contraceptive.

[0029.] It has been found that a 22 kD sperm protein, SP22, formerly identified as SP-16, is very significantly correlated with fertility ($p < 0.0001$; $N = 52$) and predictive of fertility.

If an antibody to this protein is generated, the degree of antibody recognition by sperm in an epididymal sperm sample or ejaculate would reflect the fertility of the sample. This antibody can be used as a contraceptive to block the expression of SP22 to render sperm infertile, screen for toxicity, select superior sires for artificial insemination, and select men for assisted reproductive technologies. The protein itself can be used as a contraceptive vaccine, inactivated by gene knockout, or the protein can be added to sperm from infertile men to make fertility techniques more feasible.

[0030.] SP22 is unique compared to any putative fertility proteins previously identified in either sperm or seminal plasma. Moreover, a search of the gene database failed to reveal any previously identified protein in male reproductive biology that has any homology with SP22. Both the amino acid and nucleotide sequence data indicated that SP22 is highly

09752544T525Z50

homologous with DJ-1 (Nagakubo et al., 1997), a protein for which an oncogene role was speculated. These investigators have since abandoned this notion, and, to date, no definitive role exists for DJ-1.

[0031.] The protein SP22, based upon initial electrophoretic runs, was originally thought to be a 16kD protein, with a pI of 5.5, and was so identified in great-grandparent application Serial No. 08/593,677. However, after comparing several types of molecular weight standards, the molecular weight of this protein was found consistently to have an apparent molecular weight of 22 kD in 11% acrylamide gels. For purposes of the present invention, the protein will be identified as SP22, although in the great-grandparent application USSN 08/593,677, the protein was identified as SP-16.

BRIEF DESCRIPTION OF THE DRAWINGS

[0032.] Figure 1 compares the amino acid sequence of SP22 and DJ-1, and illustrates the four peptides identified following Edman degradation that were used to identify the homology with DJ1 and correct sequences of these peptides as they exist in SP22.

[0033.] Figure 2 shows SEQ ID NO:3 which compares the nucleotide and amino acid sequences of the two SP22 mRNA transcripts, A and B. The A transcript is SP22A and is unique to the testis and referred to as SEQ ID NO:2.

[0034.] Figure 3 shows a Northern blot showing the distinct higher (approximately 1.5 kB) molecular weight band of the SEQ

097324-03030

ID NO:2 SP22A mRNA in the testis.

[0035.] Figure 4 shows silver stained two dimensional gels and corresponding immunoblots showing localization of SP22 using anti-SP22 peptide Ig in rat cauda epididymal sperm extract (top), rat cauda sperm membrane preparation (middle), and rat rete testis sperm extract (bottom).

[0036.] Figure 5 shows silver stained two dimensional gels and corresponding immunoblots showing localization of SP22 using anti-SP22 peptide Ig in detergent extracts of sperm from various species.

[0037.] Figure 6 graphs show the relationship of fertility vs. SP22 following exposure to epididymal toxicants (top) and testicular toxicants (bottom). It should be noted that the biphasic, threshold relationship is maintained regardless of the type of insult.

[0038.] Figure 7 shows data from 131 animals representing the association between SP22 and fertility. These data were pooled from a number of toxicology studies, in which testicular toxicants or epididymal toxicants were tested.

a21 [0039.] Figure 8 is a plot illustrating the immunoreactivity of overlapping 15 mer SP22 peptides with affinity-purified anti-SP22 peptide Ig. Antiserum was affinity-purified and diluted 1:100 prior to use. The two reactive peaks within the 189 amino acid SP22 sequence are: Peptide A (TVAGLAGKDPVQCSR) and Peptide B (DGLILTSR).

[0040.] Figures 9A-9C are micrographs depicting (A)

immunostaining over the equatorial segment of the head of fresh, unfixed cauda epididymal rat sperm using affinity-purified anti-SP22 peptide Ig; (B) This immunostaining is completely ablated by coincubation with 20 micrograms (10 micrograms each) of a mixture of the 15 mer and 8 mer peptides used as immunogen; (C) Immunostaining over the head of human sperm with the anti-SP22 peptide Ig; staining is restricted to the equatorial segment using anti-recombinant SP22 Ig.

[0041.] Figures 10A-C are micrographs depicting immunolocalization of SP22 on fresh, unfixed rat cauda epididymal sperm (A) in the presence of anti-SP22 peptide Ig only (1:200); (B) in the presence of anti-SP22 peptide Ig + Peptide A (20 micrograms); and (C) in the presence of anti-SP22 peptide Ig + Peptide B (20 micrograms). It should be noted that the 15 amino acid sequence of Peptide A competes effectively with the Ig, indicating that Peptide A is an exposed domain.

[0042.] Figure 11 is a graph of fertility data using affinity-purified anti-SP22 peptide Ig. When rat cauda epididymal sperm were inseminated without pre-incubation with Ig, fertility as expressed as the number of fetal implants relative to the number of corpora lutea on day 9 of gestation averaged 83%. In contrast, when sperm were incubated for five minutes with 10 microliters of affinity purified anti-SP22 peptide Ig (1:50) prior to insemination *in utero*, fertility was decreased significantly to only 7%. Indeed, only one of

09534-0000

AB

[0044.] Figure 13 is a plot illustrating the immunoreactivity of overlapping 15 mer PS-22 peptides with affinity-purified anti-recombinant SP22 Ig. Antiserum was affinity-purified and diluted 1:100 prior to use. The three reactive peaks within the 189 amino acid SP22 sequence are: Peptide C (LEEAKTQGPYDV), Peptide D (VKEILKEQENRKGLI), and Peptide E (GFGCKVTSHPKADK).

[0045.] Figure 14 is a graph depicting both *in vivo* and *in vitro* fertility results using affinity-purified anti-recombinant SP22 Ig. When rat cauda epididymal sperm were not incubated with Ig prior to *in utero* insemination, fertility was 71%. In contrast thereto, fertility was significantly reduced to 34% when recombinant SP22 Ig (1:50) was added five minutes prior to *in utero* insemination. The ability of the

antibody to recombinant SP22 to inhibit fertility was confirmed by rat *in vitro* fertilization. Approximately 70% of the eggs were fertilized, as indicated by the presence of a sperm tail within the vitellus, after overnight incubation of cauda epididymal sperm in the absence of Ig. In contrast thereto, only 39% of the eggs were fertilized when sperm were incubated in the presence of anti-recombinant SP22 Ig (1:50). Interestingly, while this antibody has been used to confirm that SP22 is primarily restricted to the equatorial segment of sperm (*cf.* Klinefelter and Welch, 1999), the distinguishing difference appears to be a significant reduction in the number of sperm binding to the zona pellucida of the egg.

[0046.] Figure 15 shows the nucleotide and amino acid sequences of SP22.

[0047.] Figure 16 shows the longer 5' untranslated regions of SP22A.

DETAILED DESCRIPTION OF THE INVENTION

[0048.] SP22 is characterized by the nucleotide and amino acid sequences of Figure 15.

[0049.] As noted previously, a number of proteins associated with the sperm membrane, including PH-20, ZP3, SP-17, and proteins D and E, as well as the Wei 17 kD protein, do not localize. In contrast, antibody to SP22 peptides localizes over a very discrete region of the sperm head, i.e., the ventral anterior surface of the equatorial segment. More importantly, however, the 17 kD protein described by Wei et

095540495260

al., (1994) was never identified in the profile of proteins in the original detergent extract. Thus, it is impossible to determine the relationship of this protein relative to others reported in the literature. It seems that this may be similar to the more recently-described SP-17 (Lea et al., 1996), but this is impossible to determine, as amino acid sequence data were not provided by Wei et al. (1994).

[0050.] It is clearly evident that SP22 is unique compared to any putative fertility proteins previously identified in either sperm or seminal plasma. Moreover, a search of the gene database failed to reveal any previously identified protein in male reproductive biology that has any homology with SP22.

[0051.] The presence of SP22 sequences with divergent 5' ends was not unexpected. Northern blotting of rat tissue RNAs, including testis, detected a 1 kB mRNA in all tissues and an additional 1.5 kB transcript found only in the testis. While a specific tissue of cell type has not been assigned to each sequence at this time, the unique 5' untranslated region of SP22 shows some similarity to the 5' untranslated region of mouse somatic expressed tag sequences homologous with rat SP22. Similarly, the longer 5' untranslated regions of SP22A, shown in Figures 2 and 16, suggested that encodes the 1.5 kB mRNA. This unique untranslated sequence may serve to impart mRNA stability for the subsequent expression of SP22 in the testis. Northern blotting with sequence specific probes is

needed to relate the different SP22 sequences to their respective mRNAs and to their relative tissue abundance.

[0052.] A short five-day exposure paradigm and multiple epididymal toxicants were used in an initial study which identified SP22 (Klinefelter et al., 1997). However, the discovery that SP22 originates in the testis prompted a study in which animals were exposed for 14 days to a testicular toxicant, bromochloroacetic acid, which is a by-product of drinking water disinfection on which the U.S. Environmental Protection Agency has requested studies, as it is one of the more prevalent disinfection by-products in drinking water. Previous studies on dibromoacetic acid (Linder et al., 1995, 1997) and dichloroacetic acid (linder et al., 1997) revealed that the di-substituted haloacetic acids perturbed spermatogenesis, and that within fourteen days, defects (i.e., alterations in sperm motion and morphology) were manifest in epididymal sperm. Therefore, it was hypothesized that bromochloroacetic acid would act similarly.

[0053.] SP22 levels on sperm were not evaluated in early haloacetic acid studies. Both a quantitative evaluation of SP22 in extracts of epididymal sperm and fertility following *in utero* insemination was incorporated in a recent study of bromochloroacetic acid. It was observed that SP22 levels were diminished in detergent extracts of epididymal sperm in a dose-related manner, with significance achieved at even the lowest dosage. Moreover, the fertility of sperm from the

0052440000

[0054.] Thus, it has now successfully been demonstrated that SP22 levels on epididymal sperm are compromised by chemicals which compromise both testicular and epididymal function. This established the feasibility of and an SP22-based assay of epididymal and ejaculated sperm as a diagnostic indicator of compromised sperm quality, i.e., fertility, in either toxicological or epidemiological settings. Additionally, the existence of SP22 on ejaculated sperm from multiple species (i.e., bull, stallion, human) established the feasibility of using such an SP22-based diagnostic to evaluate the fertility of sperm from these species when artificial breeding, herd sires, and assisted reproductive technologies (*in vitro* fertilization vs. *in utero* insemination) are considered. This is also of particular importance in breeding endangered species.

[0055.] As indicated above during discussion of the 95 kD phosphotyrosine protein discovered on the mouse sperm membrane, specific peptides inherent to this protein were able to block the binding of sperm to the zona pellucida of the egg. Presumably, these peptides competed for binding of native sperm protein. More recently, Amann et al. (1998a, b) have used small peptides comprising the saposin subunits of the prosaposin or SGP-1 protein to enhance binding of sperm to an egg membrane substrate. In this assay, increased sperm

binding indicated increased fertilizing ability.

[0056.] In the present studies, highly specific SP22 IgG was used to negate the fertility of sperm *in vivo*. Since the epitope recognized by this IgG represents 15 and 8 amino acid peptide sequences of SP22, it can be reasoned that similar site-directed, small molecule recognition has been achieved. Moreover, it seems that one or both of these peptide targets may be pivotal in the function of SP22. Thus, these and other site-directed antagonists or agonists might serve to modulate (i.e., abate or enhance) the fertility of sperm.

[0057.] The sperm protein SP22 is completely novel to the field of reproductive biology. SP22 is synthesized in the testis and can be recovered from testicular sperm before they enter the epididymis. SP22 is an integral component of the sperm membrane, and is a component of ejaculated sperm from other species including bull, stallion, and human. SP22 is highly correlated with fertility following exposure to testicular toxicants as well as epididymal toxicants.

Moreover, SP22 is a causal modulator of fertility, as anti-SP22 peptide IgG can effectively block the fertility of sperm.

[0058.] Since characteristic sequences of SP22 are known, it is possible to prepare functional derivatives of SP22 as well. By "functional derivative" is meant a fragment, variant, analog, agonist, or chemical derivative of SP22, which terms are defined below.

[0059.] A "functional derivative" retains at least a portion

of the amino acid sequence of SP22, which permits its utility in accordance with the present invention, namely, determining or affecting male fertility. A "fragment" of SP22 refers to any subset of the SP22 molecule, that is, a shorter peptide. The fragments of interest are those which can be used to determine or affect male fertility.

[0060.] A "variant" of SP22 refers to a molecule which is substantially similar to either the entire SP22 protein or fragment thereof. Variant peptides may be covalently prepared by direct chemical synthesis of the variant peptide, using methods well known in the art.

[0061.] Alternatively, amino acid sequence variants of SP22 can be prepared by mutation in the DNAs which encode the synthesized SP22. Such variants include, for example, deletions from, or insertions or substitutions of, residues within the amino acid sequence. Any combination of deletion, insertion, and substitution may also be made to arrive at the final constructs, provided that the final construct possesses the desired activity. Obviously, the mutations that will be made in the DNA encoding the variant peptide must not alter the reading frame and preferably will not create complementary regions that could produce secondary mRNA structure (cf. European Patent Publication No. EP 75,444).

[0062.] At the generic level, these variants ordinarily are prepared by site-directed mutagenesis, as exemplified by Adelman et al., DNA 2: 183, 1983, of nucleotides in the DNA

09 JUL 84 04:03:01
FBI/DOJ

encoding the peptide molecule, thereby producing DNA encoding the variant, and thereafter expressing the DNA in recombinant cell culture. The variants typically exhibit the same qualitative biological activity as the nonvariant peptide.

[0063.] An "analog" of SP22 refers to a molecule which is substantially similar to either the entire molecule or a fragment thereof. The analog may be prepared by chemical synthesis.

[0064.] A "chemical derivative" of SP22 contains additional chemical moieties not normally part of the SP22 amino acid sequence. Covalent modifications of the amino acid sequence are included within the scope of this invention. Such modifications may be introduced into the SP22 by reacting targeted amino acid residues of the peptide with an organic derivatizing agent that is capable of reacting with selected side chains or terminal residues.

[0065.] Amino terminal residues can be reacted with succinic or other carboxylic acid anhydrides. Other suitable reagents for derivatizing alpha-amino-containing residues include amidoesters such as methyl picolinimidate; pyridoxal phosphate; pyridoxal; chloroborohydrate; trinitrobenzenesulfonic acid; O-methylisourea; 2,4-pentanedione; and transaminase-catalase reacted with glyoxylate.

[0066.] Specific modifications of tyrosyl residues per se have been studied extensively, with particular interest in

introducing spectral labels into tyrosyl residues by reaction with aromatic diazonium compounds or tetranitromethane. Most commonly, N-acetylimidazole and tetranitromethane are used to form O-acetyl tyrosyl species and 3-nitro derivatives, respectively.

[0067.] Carboxyl side groups such as aspartyl or glutamyl are selectively modified by reaction with carbodiimides ($R'N-C-N-R'$) such as 1-cyclohexy-3-[2-morpholinyl-(4-ethyl)] carbodiimide or 1-ethyl-3-(4-azonia-4,4-dimethylpentyl) carbodiimide. Furthermore, aspartyl and glutamyl residues are converted to asparaginyl and glutaminyl residues by reaction with ammonium ions.

[0068.] As used herein, the term "muteins" or "variants" refers to analogs of SP22 in which one or more of the amino acid residues of the natural SP22, preferably 1-10, and more preferably 1-5, residues, or even only a single residue, are replaced by different amino acid residues or are deleted, or one or more amino acid residues, such as 1-10, 1-5, or only one residue are added to the natural sequence of SP22. These muteins are prepared by known synthesis techniques and/or site-directed mutagenesis techniques, or by any other known technique suitable therefor. The substitutions are preferably conservative, see, e.g., Schulz et al., Principle of Protein Structure, Springer-Verlag, New York, 1978; and Creighton, Proteins: Structure and Molecular Properties, W.H. Freeman & Co., San Francisco, 1983; both of which are hereby

[0069.] The types of such substitutions which may be made in the protein or peptide molecules of the present invention may be based on analysis of the frequencies of amino acid changes between a homologous protein of different species, such as those presented in Table 1-2 of Schulz et al., *op. cit.*, and Figure 3-9 of Creighton, *op.cit.* Based upon such analysis, conservative substitutions may be defined herein as exchanges within one of the following five groups:

I. Small aliphatic, nonpolar or slightly polar

Ala, Ser, Thr, Pro, Gly

II. Polar, negatively charged residues and their amides:

Asp, Asn, Glu, Gln

III. Polar, positively charged residues:

His, Arg., Lys

IV. Large, aliphatic nonpolar residues:

Met, Leu, Ile, Val, Cys

V. Large aromatic residues:

Phe, Try, Trp

[0070.] Within the foregoing groups the following five substitutions are considered "highly conservative":

Asp/Glu

His/Arg/Lys

Phe/Tyr/Trp

Met/Leu/Ile/Val

[0071.] Semi-conservative substitutions are defined to be exchanges between two of groups (I)-(V) above which are limited to supergroup (A), comprising (I), (II), and (III) above, or to supergroup (B), comprising (IV) and (V) above. Substitutions are not limited to the genetically encoded, or even the naturally occurring amino acids. When the epitope is prepared by peptide synthesis, the desired amino acid may be used directly. Alternatively, a genetically encoded amino acid may be modified by reacting it with an organic derivatizing agent that is capable of reacting with selected side chains or terminal residues.

Functional Fragments

[0072.] Functional fragments of the SP22 molecule can be identified using a three-tiered analysis, the first of which is identification of linear epitopes by mimotope analysis. For this, the 189 amino acids comprising SP22 were subdivided into 59 overlapping 15 amino acid peptides, with each peptide overlapping by three amino acids. The 59 peptides were biotinylated and allowed to bind to streptavidin-coated wells in 96-well plates. The reactivity of various antisera was detected by enzyme-linked immunosorbent assay (ELISA). Briefly, after care was taken to block non-specific binding, SP22 antibody was added sequentially (*i.e.*, either affinity-purified anti-peptide Ig or affinity-purified full-length recombinant SP22 Ig), was added followed by sequential addition of peroxidase-conjugated secondary antibody, and

peroxidase substrate. The optical density of each well was read at 450 nm and duplicate wells were averaged. The average value obtained from a similar ELISA using control serum (*i.e.*, preimmune serum) was subtracted from the test Ig values and the resultant values were plotted to determine which linear epitopes were recognized by the Ig (*cf.* Figures 9 and 13, which represent the mimotope analysis for the anti-peptide Ig and recombinant SP22 Ig, respectively).

[0073.] The second and third components in the strategy to identify functional fragments of SP22 rely on the synthesis of non-biotinylated peptides corresponding to the epitopes (peptides) predicted by the mimotope analysis. To determine whether any of the epitopes recognized by mimotope analysis are actually exposed on the surface of the sperm membrane, *i.e.*, represents an exposed domain, immunocytochemical staining with the Ig, without and with each of the peptides, was performed. For the anti-peptide Ig, immunostaining was completely ablated when peptide A was coincubated with the Ig. However, when peptide B was incubated with the Ig, staining remained identical to the staining observed with the Ig alone, as shown in Figure 10. Since we know from the mimotope analysis that the anti-peptide Ig recognizes both peptides, one can conclude that the 15 amino acid sequence of peptide A is an exposed domain, while the 8 amino acid sequence of peptide B is not exposed.

[0074.] The third component of the strategy is predicated on

00734 0074 0075 0076 0077 0078 0079 0080 0081 0082 0083 0084 0085 0086 0087 0088 0089 0090 0091 0092 0093 0094 0095 0096 0097 0098 0099 0100 0101 0102 0103 0104 0105 0106 0107 0108 0109 0110 0111 0112 0113 0114 0115 0116 0117 0118 0119 0120 0121 0122 0123 0124 0125 0126 0127 0128 0129 0130 0131 0132 0133 0134 0135 0136 0137 0138 0139 0140 0141 0142 0143 0144 0145 0146 0147 0148 0149 0150 0151 0152 0153 0154 0155 0156 0157 0158 0159 0160 0161 0162 0163 0164 0165 0166 0167 0168 0169 0170 0171 0172 0173 0174 0175 0176 0177 0178 0179 0180 0181 0182 0183 0184 0185 0186 0187 0188 0189 0190 0191 0192 0193 0194 0195 0196 0197 0198 0199 0200 0201 0202 0203 0204 0205 0206 0207 0208 0209 0210 0211 0212 0213 0214 0215 0216 0217 0218 0219 0220 0221 0222 0223 0224 0225 0226 0227 0228 0229 0230 0231 0232 0233 0234 0235 0236 0237 0238 0239 0240 0241 0242 0243 0244 0245 0246 0247 0248 0249 0250 0251 0252 0253 0254 0255 0256 0257 0258 0259 0260 0261 0262 0263 0264 0265 0266 0267 0268 0269 0270 0271 0272 0273 0274 0275 0276 0277 0278 0279 0280 0281 0282 0283 0284 0285 0286 0287 0288 0289 0290 0291 0292 0293 0294 0295 0296 0297 0298 0299 0300 0301 0302 0303 0304 0305 0306 0307 0308 0309 0310 0311 0312 0313 0314 0315 0316 0317 0318 0319 0320 0321 0322 0323 0324 0325 0326 0327 0328 0329 0330 0331 0332 0333 0334 0335 0336 0337 0338 0339 0340 0341 0342 0343 0344 0345 0346 0347 0348 0349 0350 0351 0352 0353 0354 0355 0356 0357 0358 0359 0360 0361 0362 0363 0364 0365 0366 0367 0368 0369 0370 0371 0372 0373 0374 0375 0376 0377 0378 0379 0380 0381 0382 0383 0384 0385 0386 0387 0388 0389 0390 0391 0392 0393 0394 0395 0396 0397 0398 0399 0400 0401 0402 0403 0404 0405 0406 0407 0408 0409 0410 0411 0412 0413 0414 0415 0416 0417 0418 0419 0420 0421 0422 0423 0424 0425 0426 0427 0428 0429 0430 0431 0432 0433 0434 0435 0436 0437 0438 0439 0440 0441 0442 0443 0444 0445 0446 0447 0448 0449 0450 0451 0452 0453 0454 0455 0456 0457 0458 0459 0460 0461 0462 0463 0464 0465 0466 0467 0468 0469 0470 0471 0472 0473 0474 0475 0476 0477 0478 0479 0480 0481 0482 0483 0484 0485 0486 0487 0488 0489 0490 0491 0492 0493 0494 0495 0496 0497 0498 0499 0500 0501 0502 0503 0504 0505 0506 0507 0508 0509 0510 0511 0512 0513 0514 0515 0516 0517 0518 0519 0520 0521 0522 0523 0524 0525 0526 0527 0528 0529 0530 0531 0532 0533 0534 0535 0536 0537 0538 0539 0540 0541 0542 0543 0544 0545 0546 0547 0548 0549 0550 0551 0552 0553 0554 0555 0556 0557 0558 0559 0560 0561 0562 0563 0564 0565 0566 0567 0568 0569 0570 0571 0572 0573 0574 0575 0576 0577 0578 0579 0580 0581 0582 0583 0584 0585 0586 0587 0588 0589 0590 0591 0592 0593 0594 0595 0596 0597 0598 0599 0600 0601 0602 0603 0604 0605 0606 0607 0608 0609 0610 0611 0612 0613 0614 0615 0616 0617 0618 0619 0620 0621 0622 0623 0624 0625 0626 0627 0628 0629 0630 0631 0632 0633 0634 0635 0636 0637 0638 0639 0640 0641 0642 0643 0644 0645 0646 0647 0648 0649 0650 0651 0652 0653 0654 0655 0656 0657 0658 0659 0660 0661 0662 0663 0664 0665 0666 0667 0668 0669 0670 0671 0672 0673 0674 0675 0676 0677 0678 0679 0680 0681 0682 0683 0684 0685 0686 0687 0688 0689 0690 0691 0692 0693 0694 0695 0696 0697 0698 0699 0700 0701 0702 0703 0704 0705 0706 0707 0708 0709 0710 0711 0712 0713 0714 0715 0716 0717 0718 0719 0720 0721 0722 0723 0724 0725 0726 0727 0728 0729 0730 0731 0732 0733 0734 0735 0736 0737 0738 0739 0740 0741 0742 0743 0744 0745 0746 0747 0748 0749 0750 0751 0752 0753 0754 0755 0756 0757 0758 0759 0760 0761 0762 0763 0764 0765 0766 0767 0768 0769 0770 0771 0772 0773 0774 0775 0776 0777 0778 0779 0780 0781 0782 0783 0784 0785 0786 0787 0788 0789 0790 0791 0792 0793 0794 0795 0796 0797 0798 0799 0800 0801 0802 0803 0804 0805 0806 0807 0808 0809 0810 0811 0812 0813 0814 0815 0816 0817 0818 0819 0820 0821 0822 0823 0824 0825 0826 0827 0828 0829 0830 0831 0832 0833 0834 0835 0836 0837 0838 0839 0840 0841 0842 0843 0844 0845 0846 0847 0848 0849 0850 0851 0852 0853 0854 0855 0856 0857 0858 0859 0860 0861 0862 0863 0864 0865 0866 0867 0868 0869 0870 0871 0872 0873 0874 0875 0876 0877 0878 0879 0880 0881 0882 0883 0884 0885 0886 0887 0888 0889 0890 0891 0892 0893 0894 0895 0896 0897 0898 0899 0900 0901 0902 0903 0904 0905 0906 0907 0908 0909 0910 0911 0912 0913 0914 0915 0916 0917 0918 0919 0920 0921 0922 0923 0924 0925 0926 0927 0928 0929 0930 0931 0932 0933 0934 0935 0936 0937 0938 0939 0940 0941 0942 0943 0944 0945 0946 0947 0948 0949 0950 0951 0952 0953 0954 0955 0956 0957 0958 0959 0960 0961 0962 0963 0964 0965 0966 0967 0968 0969 0970 0971 0972 0973 0974 0975 0976 0977 0978 0979 0980 0981 0982 0983 0984 0985 0986 0987 0988 0989 0990 0991 0992 0993 0994 0995 0996 0997 0998 0999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050 1051 1052 1053 1054 1055 1056 1057 1058 1059 1060 1061 1062 1063 1064 1065 1066 1067 1068 1069 1070 1071 1072 1073 1074 1075 1076 1077 1078 1079 1080 1081 1082 1083 1084 1085 1086 1087 1088 1089 1090 1091 1092 1093 1094 1095 1096 1097 1098 1099 1100 1101 1102 1103 1104 1105 1106 1107 1108 1109 1110 1111 1112 1113 1114 1115 1116 1117 1118 1119 1120 1121 1122 1123 1124 1125 1126 1127 1128 1129 1130 1131 1132 1133 1134 1135 1136 1137 1138 1139 1140 1141 1142 1143 1144 1145 1146 1147 1148 1149 1150 1151 1152 1153 1154 1155 1156 1157 1158 1159 1160 1161 1162 1163 1164 1165 1166 1167 1168 1169 1170 1171 1172 1173 1174 1175 1176 1177 1178 1179 1180 1181 1182 1183 1184 1185 1186 1187 1188 1189 1190 1191 1192 1193 1194 1195 1196 1197 1198 1199 1200 1201 1202 1203 1204 1205 1206 1207 1208 1209 1210 1211 1212 1213 1214 1215 1216 1217 1218 1219 1220 1221 1222 1223 1224 1225 1226 1227 1228 1229 1230 1231 1232 1233 1234 1235 1236 1237 1238 1239 1240 1241 1242 1243 1244 1245 1246 1247 1248 1249 1250 1251 1252 1253 1254 1255 1256 1257 1258 1259 1260 1261 1262 1263 1264 1265 1266 1267 1268 1269 1270 1271 1272 1273 1274 1275 1276 1277 1278 1279 1280 1281 1282 1283 1284 1285 1286 1287 1288 1289 1290 1291 1292 1293 1294 1295 1296 1297 1298 1299 1300 1301 1302 1303 1304 1305 1306 1307 1308 1309 1310 1311 1312 1313 1314 1315 1316 1317 1318 1319 1320 1321 1322 1323 1324 1325 1326 1327 1328 1329 1330 1331 1332 1333 1334 1335 1336 1337 1338 1339 1340 1341 1342 1343 1344 1345 1346 1347 1348 1349 1350 1351 1352 1353 1354 1355 1356 1357 1358 1359 1360 1361 1362 1363 1364 1365 1366 1367 1368 1369 1370 1371 1372 1373 1374 1375 1376 1377 1378 1379 1380 1381 1382 1383 1384 1385 1386 1387 1388 1389 1390 1391 1392 1393 1394 1395 1396 1397 1398 1399 1400 1401 1402 1403 1404 1405 1406 1407 1408 1409 1410 1411 1412 1413 1414 1415 1416 1417 1418 1419 1420 1421 1422 1423 1424 1425 1426 1427 1428 1429 1430 1431 1432 1433 1434 1435 1436 1437 1438 1439 1440 1441 1442 1443 1444 1445 1446 1447 1448 1449 1450 1451 1452 1453 1454 1455 1456 1457 1458 1459 1460 1461 1462 1463 1464 1465 1466 1467 1468 1469 1470 1471 1472 1473 1474 1475 1476 1477 1478 1479 1480 1481 1482 1483 1484 1485 1486 1487 1488 1489 1490 1491 1492 1493 1494 1495 1496 1497 1498 1499 1500 1501 1502 1503 1504 1505 1506 1507 1508 1509 1510 1511 1512 1513 1514 1515 1516 1517 1518 1519 1520 1521 1522 1523 1524 1525 1526 1527 1528 1529 1530 1531 1532 1533 1534 1535 1536 1537 1538 1539 1540 1541 1542 1543 1544 1545 1546 1547 1548 1549 1550 1551 1552 1553 1554 1555 1556 1557 1558 1559 1560 1561 1562 1563 1564 1565 1566 1567 1568 1569 1570 1571 1572 1573 1574 1575 1576 1577 1578 1579 1580 1581 1582 1583 1584 1585 1586 1587 1588 1589 1590 1591 1592 1593 1594 1595 1596 1597 1598 1599 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615 1616 1617 1618 1619 1620 1621 1622 1623 1624 1625 1626 1627 1628 1629 1630 1631 1632 1633 1634 1635 1636 1637 1638 1639 1640 1641 1642 1643 1644 1645 1646 1647 1648 1649 1650 1651 1652 1653 1654 1655 1656 1657 1658 1659 1660 1661 1662 1663 1664 1665 1666 1667 1668 1669 1670 1671 1672 1673 1674 1675 1676 1677 1678 1679 1680 1681 1682 1683 1684 1685 1686 1687 1688 1689 1690 1691 1692 1693 1694 1695 1696 1697 1698 1699 1700 1701 1702 1703 1704 1705 1706 1707 1708 1709 1710 1711 1712 1713 1714 1715 1716 1717 1718 1719 1720 1721 1722 1723 1724 1725 1726 1727 1728 1729 1730 1731 1732 1733 1734 1735 1736 1737 1738 1739 1740 1741 1742 1743 1744 1745 1746 1747 1748 1749 1750 1751 1752 1753 1754 1755 1756 1757 1758 1759 1760 1761 1762 1763 1764 1765 1766 1767 1768 1769 1770 1771 1772 1773 1774 1775 1776 1777 1778 1779 1780 1781 1782 1783 1784 1785 1786 1787 1788 1789 1790 1791 1792 1793 1794 1795 1796 1797 1798 1799 1800 1801 1802 1803 1804 1805 1806 1807 1808 1809 1810 1811 1812 1813 1814 1815 1816 1817 1818 1819 1820 1821 1822 1823 1824 1825 1826 1827 1828 1829 1830 1831 1832 1833 1834 1835 1836 1837 1838 1839 1840 1841 1842 1843 1844 1845 1846 1847 1848 1849 1850 1851 1852 1853 1854 1855 1856 1857 1858 1859 1860 1861 1862 1863 1864 1865 1866 1867 1868 1869 1870 1871 1872 1873 1874 1875 1876 1877 1878 1879 1880 1881 1882 1883 1884 1885 1886 1887 1888 1889 1890 1891 1892 1893 1894 1895 1896 1897 1898 1899 1900 1901 1902 1903 1904 1905 1906 1907 1908 1909 1910 1911 1912 1913 1914 1915 1916 1917 1918 1919 1920 1921 1922 1923 1924 1925 1926 1927 1928 1929 1930 1931 1932 1933 1934 1935 1936 1937 1938 1939 1940 1941 1942 1943 1944 1945 1946 1947 1948 1949 1950 1951 1952 1953 1954 1955 1956 1957 1958 1959 1960 1961 1962 1963 1964 1965 1966 1967 1968 1969 1970 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980 1981 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2

the hypothesis that only epitopes having exposed domains play functional roles in fertility. To test this, cauda epididymal sperm were inseminated *in utero* following incubation with Ig alone, or incubation with Ig and individual peptides (*i.e.*, peptide A and peptide B). Indeed, while the anti-peptide Ig almost completely inhibited fertility, coincubation of Ig and peptide A resulted in no alteration in fertility relative to historical values. In contrast therewith, coincubation of anti-peptide Ig and peptide B also resulted in near complete inhibition of fertility, as shown in Figure 12.

[0075.] Collectively, these data clearly demonstrate that the exposed 15 amino acids of Peptide A represent a functional fragment of the SP22 molecule. In a similar fashion, any peptide can be assayed to determine if it, too, is a functional fragment of SP22.

[0076.] Based on work with Ig to the full length recombinant SP22, it can be reasoned that other functional fragments exist. First, the mimotope analysis of the recombinant SP22 Ig shown in Figure 13 reveals three peptide epitopes. Second, the Ig to the recombinant SP22 immunostains sperm of all species tested (rat, bull, rabbit, hamster, and human; data for rat, bull, and human have been shown by Klinefelter and Welch, 2000). Third, this Ig inhibits fertility both *in vivo* and *in vitro* as shown in Figure 14. As indicated above, any fragments can readily be tested without undue experimentation to identify those fragments which modulate fertility.

Targeting functional fragments is likely to effectively reduce fertility to 0% and makes the use of site-directed antagonists more feasible across many species, including humans.

[0077.] Antibodies that identify functional SP22 fragments also lend themselves to testing in random peptide phase display technology (Eidne et al., 2000). Briefly, fifteen or twelve-mer random peptide phase display libraries can be used to determine what peptides might interact with functional SP22 peptides by competitive displacement of Fab fragments of SP22 antibodies. For this, fixed sperm are allowed to adhere to wells in multiwell plates, and immunostaining for SP22 is evaluated in the absence and presence of unique and random peptides expressed by the phage library. Once the competitive peptides are identified by amino acid sequence analysis, increased amounts of peptide can be synthesized and used as alternative molecular antagonists to antibodies directed against functional fragments.

[0078.] Another alternative is to screen thousands of small molecule organics for their ability to competitively display Fab fragments to functional SP22 fragments.

Screening with SP22 Antibody

[0079.] It is apparent from the above description of SP22 antibodies that a wide variety of diagnostic tests is possible using the antibodies of the invention. In attempting to diagnose causes of infertility, an immunoassay to detect decreased levels of SP22 on sperm is a useful adjunct to known

007774 007774

hormone assay. Further uses of the antibodies include testing livestock for candidates for artificial insemination: the higher the levels of SP22 in the potential donor, the more likely artificial insemination is to be successful. Isolation of SP22 allows production of an antisera containing antibody to SP22 for possible crossreaction with other species, including human SP22. This antibody enables preparation of an enzyme-linked immunosorbent assay (ELISA).

[0080.] For example, to evaluate antibody binding, polystyrene microwells were precoated with extract of a particular epididymal sperm (rat) or ejaculate (horse, bull, human) sample containing an unknown amount of SP22. Next, SP22 antibody was added, followed by the addition of avidin-biotin-peroxidase complex. A precipitate formed when a substrate such as DAB is oxidized by peroxidase in the presence of hydrogen peroxide. A standard curve for SP22 was generated using increasing known amounts of SP22. The amount of SP22 in a sample is then determined by the optical density of the colored precipitate in the sample and the linear regression obtained from the set of SP22 standards.

[0081.] Aside from ELISA, the amount of SP22 present on the surface of sperm in a sample (epididymal or ejaculate, animal or human) can be determined using quantitative fluorescence spectroscopy or fluorescent light microscopy. For this, sperm are incubated with SP22 antibody and then with labelled Rhodamine or FITC-conjugated second antibody. It is first

necessary to determine the relationship between fluorescence of a sample in a fluorometer or a microscopic image, and the optical density of SP22 separated by two dimensional gel electrophoresis. Once this is established, fluorescence can be related to fertility.

[0082.] It is also important to determine the relationship between the number of sperm in a sample which express (SP22) (i.e., fluoresce), the degree of the expression or fluorescence, and fertility. This is particularly true for men considering assisted reproductive technologies. For example, if only a critical number (X) of sperm is needed to express a threshold amount (Y) of SP22 for a successful attempt at fertility, it is possible to selectively remove those sperm not expressing SP22 in the ejaculate and use only those sperm that do express a sufficient amount of SP22, for assisted reproductive technologies such as intra uterine transfer or IVF following dissociation of SP22 expressing sperm from the SP22 antibody.

[0083.] To determine if there is a relationship between the number of sperm expressing SP22 and the extent to which they express it, sperm binding SP22 antibody or antagonist are evaluated by quantitative indirect fluorescence microscopy. For this, Rhodamine or FITC immunolabeling is performed on an aliquot of sperm equivalent to that used for *in utero* insemination, and the number of sperm that fluoresce is determined along with the relative degree of fluorescence of

individual sperm in a sample. The resulting fluorescence histograms are related to fertility assessed by artificial (*in utero*) insemination. To determine whether a critical number of SP22 expressing sperm are requisite to fertility, an aliquot equivalent to that used for *in utero* insemination is subjected to immunoabsorption. Polystyrene microwells are precoated with SP22 antibody and sperm in the ejaculate are allowed to bind. Those sperm not binding are washed away, antibody-bound sperm are recovered following dissociation of the antibody with incubation in 0.1 M lithium diiodosalicylate, and increasing numbers of these SP-22 expressing sperm are inseminated in utero.

[0084.] Conservative amino acid substitutions according to the present invention, e.g., as presented above, are known in the art and would be expected to maintain the biological and structural properties of the polypeptide after such amino acid substitutions. Most deletions, insertions, and substitutions according to the present invention are those which do not produce radical changes in the characteristics of the protein or peptide molecules. One skilled in the art will appreciate that the effect of substitution can be evaluated by routine screening assays, either immunoassays or bioassays. For example, a mutant typically is made by site-specific mutagenesis of the peptide molecule-encoding nucleic acid, expression of the mutant nucleic acid in recombinant cell culture, and, optionally, purification from the cell culture,

0075254-010394

or a biological sample containing SP22, for example, by immunoaffinity chromatography using a specific antibody on a column to absorb the mutant by binding to at least one epitope.

SP22 Amino Acid Sequencing

[0085.] Cauda epididymal sperm were extract for one hour as described previously (Klinefelter et al., 1997) using 80 mM n-octyl- β -glucopyranoside (OBG) in 10 mM Tris, pH 7.2, to which 0.2 mM phenylmethanesulfonyl fluoride was freshly added. The extract was then concentrated, desalted, and assayed for protein prior to HPLC separation. Fractions highly enriched in SP22 were obtained by reverse phase C4 HPLC using a linear gradient of 20-80% acetonitrile in water with 0.1% TFA. SP22-containing fractions were pooled and aliquots equivalent to 60 μ g of protein were loaded for separation on two dimensional 14% SDS-PAGE. Gels were then stained with Coomassie Blue and the SP22 spots punched out and frozen for use in subsequent peptide sequencing. Isolated SP22 protein then was subjected to tryptic digestion, and the resulting peptide mixture was separated by HPLC. Peaks representing homogeneous peptides were selected for sequence determination by Edman degradation, and the resulting peptide sequences were matched against NCBI GenBank protein sequences using the BLAST program.

[0086.] Figure 1A shows the SP22 peptides identified by Edman degradation. Figure 1B shows the full length amino acid sequence of human DJ1. In the Figure, * indicates that one of

the threonines (T) in peptide #3 is a Serine (S) in SP22. * indicates that the initial amino acid after the lysine (K) cleavage site, i.e., valine (V), was ambiguous upon sequencing.

Methods

[0087.] Cauda epididymal sperm were extracted for one hour as described previously (Klinefelter et al., 1997) using 80 mM n-octyl- β -glucopyranoside (OBG) in 10mM Tris buffer, pH 7.2., to which 0.2 mM phenylmethanesulfonyl fluoride was freshly added. The extract was then concentrated, desalted, and assayed for protein prior to HPLC separation. Fractions highly enriched in SP22 were obtained by reverse phase C4 HPLC using a linear gradient of 20-80% acetonitrile in water with 0.1% TFA. SP22-containing fractions were pooled and aliquots equivalent to 60 μ g of protein were loaded for separation by two dimensional 14% SDS-PAGE. Gels were then stained with Coomassie Blue and the SP22 spots punched out and frozen for use in subsequent peptide sequencing. Isolated SP22 protein then was subjected to tryptic digestion and the resulting peptide mixture separated by HPLC. Peaks representing homogeneous peptides were selected for sequence determination by Edman degradation, and the resulting peptide sequences were matched against NCBI GenBank protein sequences using the BLAST program.

Sequencing SP22 cDNA and Northern Blotting

[0088.] The partial amino acids identified for SP22 were

097534-0403

substantially homologous with human DJ-1 (Nagakubo et al., 1997), and a rat testis cDNA library (Stratagene, LaJolla, CA) was screened with an EST cDNA (Accession No. AA388672) encoding a mouse DJ-1 homology. A DJ-1 cDNA probe was prepared by random primer labeling with [³²P]-dCTP (Amersham, Arlington Heights, IL) using a Prime-It II kit (Stratagene). Library screening and bacteriophage isolation was carried out using the method of Benton and Davis (1977).

[0089.] SP22 insert DNA was sequenced using the dideoxynucleotide termination method of Sanger et al., 1977, using the SequiTherm Excel Kit (Epicenter Technologies, Madison, WI). Sequence data were assembled using the MacVector analysis package (Oxford Molecular Products, Oxford, England).

[0090.] To evaluate the tissue specificity of SP22, total RNA was isolated from multiple reproductive and somatic tissues. Northern blotting to 10 µg total RNA was performed as described by Welch et al., 1992, with a stringent wash temperature of 60°C and an exposure time of 20 hours.

SP22 Peptide Antibody Location

[0091.] Peptides #1 and #4 (Figure 1) obtained by Edman degradation were used together as antigen to generate polyclonal antibody. For this, each peptide was synthesized, conjugated to carrier protein, and the two peptide-carrier protein conjugates were used to immunize two four year old female Broder Leicester Merino Sheep (service provided by

Chiron Technologies, Raleigh, NC). Specifically, each conjugate was emulsified in 1ml of Freund's complete adjuvant (approx. 0.3 mg each peptide) followed by intramuscular injection. Similar injections were administered two and six weeks later using Freund's incomplete adjuvant. Serum was collected two weeks after the final injection.

[0092.] Peptides #1 and #4 (2 mg each) were also coupled to CNBr-activated Sepharose and used for affinity purification of anti-SP22 peptide IgG. Briefly, 10 ml of immune serum were mixed with 1 ml peptide-linked Thiopropyl-Sepharose 6B overnight at 4°C. Bound anti-SP22 peptide IgG was eluted with 0.1 M glycine-HCl, pH 2.5. IgG was subsequently neutralized, desalted, concentrated, and assayed.

[0093.] Affinity-purified anti-SP22 peptide IgG (2mg/ml) was used to localize SP22 in both two dimensional gels and on sperm. For immunoblotting, proteins in sperm extracts were first resolved by mini-two dimensional gel electrophoresis, and subsequently transferred onto PVDF membranes. The blotted membranes were incubated for one hour at 34°C in DPBS with 1% BSA containing 10% normal rabbit serum. Next, affinity purified anti-SP22 peptide IgG (1:1000) was added and blots were allowed to shake overnight at 4°C. Biotinylated rabbit anti-sheep IgG and ABC reagents were added as per Vectastain instructions and the peroxides reaction product was visualized using the VIP substrate kit. To control for nonspecific binding, pre-immune serum was used in place of the affinity-

purified anti-SP22 peptide IgG.

[0094.] For immunocytochemistry, cauda epididymal (rat) or ejaculated sperm (bull, human) were washed twice with DPBS and either fixed in Zamboni's fixative containing 0.1% Triton X-100 for one hour at 4°C or incubated directly in blocking buffer (DPBS containing 1% BSA and 10% normal rabbit serum) for one hour at 34°C. Fixed sperm were incubated in blocking buffer after fixation. Blocking buffer was removed after centrifugation and 10×10^6 sperm were incubated in 1ml of Dulbecco's Phosphate Buffered Saline (DPBS) containing 20 µg of affinity-purified anti-SP22 peptide IgG (1:100) for one hour. After washing, FITC-labeled rabbit anti-sheep IgG (1:25) was added for one hour. Sperm were washed again and mounted using anti-fade mounting medium. The specificity of immunostaining was verified by adding a 20 µg mixture of peptides #1 and #4 (10 µg each) in conjunction with the affinity-purified anti-SP22 IgG.

Modulation of Fertility with SP22 Peptide Antibody

[0095.] Artificial (*in utero*) insemination in the rat was conducted as previously described (Klinefelter et al., 1997). Briefly, 10×10^6 cauda epididymal sperm were incubated for five minutes at 34°C either with or without 10µl of the affinity-purified anti-SP22 IgG (1:50), and 5×10^6 were injected into each uterine horn of LHRH-synchronized, cervically-stimulated adult females while under halothane anesthesia. Nine days later, the inseminated females were sacrificed and the number

097654-0030

of implants and corpora lutea were enumerated. Fertility was expressed as the number of implants relative to the number of corpora lutea.

Quantification of SP22 to Detect Alterations in Fertility due to Testicular Toxicants

[0096.] Bromochloroacetic acid (BCA), a naturally-occurring by-product of drinking water disinfection, was administered to adult male rats in water by gavage in graded doses, i.e., 0, 8, 24, and 72 mg/kg body weight. The rats were dosed daily for fourteen days. On day fifteen, sperm from the proximal cauda epididymis were prepared for artificial insemination as described above. The sperm remaining after insemination were washed and extracted with 80 mM n-octyl- β -glucopyranoside (OBG) in 10mM Tris, pH 7.2. The extract was then concentrated, desalted, and protein concentration was determined prior to separation on 14% mini, two-dimensional SDS-PAGE gels. The silver-stained SP22 protein was background corrected and the integrated optical density was correlated with the fertility of these sperm.

SP22 Amino Acid Sequencing

[0097.] Four peptides were obtained as shown in Figure 1. Each peptide was relatively short in length and each was flanked by trypsin cleavage sites at Lys (K) or Arg (R). Peptides #1, 2, and 4 matched sequences in the DJ-1 protein recently described in human Hela cells (Nagakubo et al., 1997). Moreover, five of the seven amino acids contained within peptide #3 following Edman degradation matched the DJ-1

09752514-010301

sequence. Of the two amino acids in peptide #3 which did not match DJ-1 sequence, one is now known to be erroneous (i.e., G should be H), and the other represents a T (human DJ-1) to S (rat SP22) substitution in the DNA sequence (Figure 2).

Comparisons of human DJ-1 with expressed sequence tagged cloned from mouse indicated that these peptides were also perfectly conserved between human and mouse.

[0098.] Figure 1 shows the SP22 peptides identified by Edman degradation and the full length amino acid sequence of SP22 and the corresponding homology to human DJ-1.

Cloning and Sequencing SP22 cDNA

[0099.] Sequencing of SP22 cDNAs obtained from a rat testis cDNA library indicated that SP22 was encoded by two distinct mRNA sequences with divergent 5' sequences (Figure 2).

[0100.] Of the four peptides derived from direct peptide sequencing of purified SP22, Figure 1, peptides #1, 2 and 4 were found to match the predicted SP22 amino acid sequence exactly. Peptide #3 was found to contain a single discrepancy where glycine was predicted rather than histidine. The 3' untranslated region of the SP22 cDNAs contained a typical polyadenylation signal (AATAAA), although separate sites of polyadenylation were observed. Database searches using the SP22 sequence indicated a substantial homology (91% identity) with the human DJ-1 protein (Figure 1) and suggest that SP22 and DJ-1 are members of the same protein family. All matching with DJ-1 was confined to the 189 amino acid residues common

00984-01001

to the three SP22 transcripts.

[0101.] Figure 2 demonstrates that SP22 is encoded by alternatively-spliced mRNAs. The divergent 5' ends of three SP22 cDNAs designated A (plain text) and B (*italics*). Peptide sequencing matched perfectly with the exception of peptide #3, where a histidine (H*) was observed in place of glycine (G). The canonical polyadenylation signal (AATAAA) is underlined. Observed multiple polyadenylation sites are indicated by asterisks.

[0102.] Figure 1 illustrates the substantial homology of rat SP22 with human DJ-1 protein. The 189 amino acids of SP22 conserved between all SP22 sequences (upper) are compared with the complete DJ-1 sequence (lower). Sequence identities are indicated by solid bars, conservative substitutions are shown by a colon, and divergent residues are indicated by gaps. All peptide sequences are identical between SP22 and DJ-1 with the exception of peptide #3, where the observed threonine (T) to serine (S) change was observed both in the directed and predicted amino acid sequences. The high degree of identity suggests that SP22 and DJ-1 belong to the same family of proteins.

[0103.] Figure 3, shows that Northern blotting of total RNA from rat testis (T), epididymis (E), brain (B), liver (L), and kidney (K) revealed the presence of a 1.0 kb mRNA in all tissues. However, a 1.5 kb transcript also appeared in the testis lane, indicating the presence of the testis-specific

0975254-040304

SP22A mRNA (SEQ ID NO: 2).

Peptide Antibody Localization

[0104.] Affinity-purified anti-SP22 peptide IgG recognized SP22 in the detergent extract of cauda epididymal sperm, solubilized membranes isolated from cauda epididymal sperm, and a detergent extract of sperm recovered from the rat testis 18 hours after efferent duct ligation (Figure 4). No signal was detected on blots incubated with preimmune serum. The fact that a slightly more basic protein at the same apparent molecular weight was also recognized by the affinity-purified anti-SP22 peptide IgG suggests that post-translationally modified variants of SP22 exist. When affinity-purified anti-SP22 peptide IgG was used to probe immunoblots of detergent extracts of bull, rabbit, stallion, and human sperm, a pattern of immunorecognition identical to that seen for the rat was evident (Figure 5), suggesting that SP22 and its isoform(s) are present in the sperm membrane regardless of species.

[0105.] Affinity-purified anti-SP22 peptide Ig localized over the anterior ventral, *i.e.*, equatorial, region of the head on either fixed or fresh, unfixed cauda epididymal rat sperm (Figure 9). Immunostaining was completely ablated by coincubating the Ig with a mixture of the immunogenic SP22 peptides, *i.e.*, peptides #1 and #4 (equivalent to peptides A and B from mimotope analyses). When peptides #1 and #4 were tested separately during coincubation with the anti-peptide Ig, the results indicated that only peptide #1 was exposed on

09254-00301
T02070-455250

fresh sperm. Coincubation with peptide #1 ablated all immunostaining, while coincubation with peptide #4 did not influence the level of immunostaining (Figure 10). Staining was also evident over the head of human sperm (Figure 9). Using the affinity-purified anti-recombinant SP22 Ig, this staining was restricted to the equatorial segment of sperm from all species examined, including human.

Modulation of Fertility with SP22 Peptide Antibody

[0106.] When cauda epididymal sperm were incubated for five minutes with anti-SP22 peptide antibody (1:50) just prior to insemination into the uterine horns of receptive females, fertility was significantly reduced (Figure 11). Indeed, while fertility of sperm that was not incubated with antibody averaged 83% (ranging from 64 to 100%), only one of the six females inseminated with sperm that were incubated with antibody had any implants. The fertility of this one female was below normal (44%).

[0107.] To repeat and extend these results, affinity-purified anti-peptide Ig (10 microliters, equivalent to 20 micrograms Ig) was incubated 1:50 (0.04 microgram/microliter) with cauda epididymal sperm five minutes prior to insemination. Fertility was once again reduced to less than 10% (Figure 12). In contrast thereto, when 20 micrograms of peptide #1, equivalent to Peptide A from mimotope analysis, was added to the Ig and sperm for five minutes prior to insemination, the resultant fertility was equivalent to

09752544-010304

Correlation of SP22 with Infertility Induced by a Testicular Toxicant

(1) establish a threshold value for the background-corrected, integrated optimal density of SP22 based on the point on the predicted line with the smallest 95% confidence

[0110.] A recombinant SP22 expression cassette encoding the entire SP22 protein was synthesized by PCR amplification of the coding region from the SP22 cDNA. This cassette was cloned into a pQE8 plasmid containing a prokaryotic lac promoter region and the Shine-Delgado ribosome binding site to facilitate expression in *E. coli*. The plasmid also contained a sequence encoding a series of six histine residues (6xHis) to allow rapid purification of the recombinant protein. Once transfected into *E. coli*, SP22 production was induced by the addition of isopropyl-thio- β -D-galactopyranoside (IPTG) to activate the lac promoter. The 6xHis tag has a pH-dependent high affinity for nickel and is capable of binding in the presence of high concentration of urea and guanidinium salts. After growth in culture for 5-6 hours, cells were harvested by centrifugation and solubilized in 6M guanidine hydrochloride, 100 mM phosphate at pH 8.0. The nickel agarose column was washed stepwise in 8M urea, 100 mM phosphate buffer at pH 8.0,

pH 6.3, pH 5.9, NS DINlly pH 4.5. The bacterial proteins were either not retained on the column or were eluted in the first three washes, while recombinant SP22 (rSP22) was eluted at pH 4.5. The purified rSP22 yields for one liter of bacterial culture were in the milligram range.

[0111.] Purified rSP22 was used unconjugated as antigen to immunize two four year old Border Leicester Merino sheep (service provided by Chrion Mimotopes, Clayton Victoria, Australia). Briefly, 1 mg of rSP22 was resuspended in 1 ml of phosphate buffered saline and emulsified with an equal volume of Complete Freund's Adjuvant and approximately 1 ml (half of the total volume) was injected into each sheep intramuscularly. A second and third immunization followed two and three weeks later, using Incomplete Freund's Adjuvant. Sera were tested by ELISA for rSP22-specific antibody titre. Sera exhibited high titre based on ELISA and anti-rSP22 Ig was purified by affinity chromatography on a Sepharose column conjugated with 25 mg of rSP22. Anti-rSP22 Ig was tested for ability to inhibit fertility both *in vivo* and *in vitro*. For *in utero* inseminations, 10 microliters (equivalent to 60 micrograms) was incubated 1:50 with rat cauda epididymal sperm, and sperm were allowed to incubate five minutes prior to insemination. For *in vitro* fertilization, a similar Ig concentration was incubated with cauda epididymal sperm in the presence of eggs overnight. Fertility was assessed *in vivo* on day 9 of gestation by the number of fetal implants relative to

the number of corpora lutea. Fertilization was assessed *in vitro* by the percentage of eggs containing a sperm tail the next morning. In addition the relative number of sperm binding to the zona after insemination was evaluated.

Antibodies and Other Molecular Antagonists to SP22

[0112.] Antibodies to SP22 can be prepared by conventional polyclonal or monoclonal technologies. These antibodies can be raised in rabbits, mice, sheep, or tissue culture cells derived therefrom, or can be products of cells of human origin.

[0113.] To prepare the antibodies, purified native SP22, SP22 peptides (*i.e.*, fragments), or recombinant SP22 (full length or functional fragments) may be used to effectively immunize animals. These antibodies may be generated either in host animals or in recipients *in vivo* (*i.e.*, via vaccine formulation). It should be noted that with respect to vaccine generation, functional fragments specific to the native molecule should be selected. Moreover, antibodies to multiple functional fragments may be generated to enhance inhibition of fertility. Antibodies generated in host animals may be administered by periodic injections (male or female), or by vaginal delivery upon emulsification in appropriate stabilizer/carrier. Similarly, peptide ligands shown to competitively inhibit binding of antibodies to functional SP22 fragments (*e.g.*, a peptide ligand for Peptide A) would be expected to effectively inhibit fertility. Again, such

09254-04034

ligands, emulsified in similar delivery systems, could be administered singularly or in combination until maximum inhibition was achieved.

Antibodies to SP22

[0114.] Antibodies to SP22 can be prepared by any conventional means, and they can be either polyclonal or monoclonal. They may be raised in rabbits, mice, or other animals or tissue culture cells derived therefrom, or can be products or cells of human origin. They may also be produced by recombinant DNA technology either in a form identical to that of the native antibody or as chimeric molecules, constructed by recombination of antibody molecules of human or animal origin or in other forms chosen to make the antibodies most suitable for use in therapy.

[0115.] For preparation of the antibodies, either purified SP22 or a peptide identical to the known sequences of fragments thereof, e.g., to the N-terminal protein sequence, may be used to immunize animals. A further possibility is to fuse one of the nucleotide sequences coding an active fragment of SP22 to the gene coding for protein A, to express the antibody. The antibody is then purified by affinity chromatography on a Sepharose column and used to immunize animals.

Preparation of Antibodies to SP22

[0116.] Antibodies (both polyclonal and monoclonal) to SP22 may be prepared for diagnostic and therapeutic uses, including

09763544-040004

but not limited to fertility control (contraception) and fertility assessment (screening).

[0117.] "Antibody" in this context refers to a synthetic protein which binds SP22 and negates its biological function. Antibodies to SP22 are prepared by either polyclonal or monoclonal techniques:

[0118.] A. Polyclonal Antibody Production

For polyclonal antibody production, adult sheep, mice or rabbits are immunized with 25 or 100 mg of SP22 suspended in Freund's complete adjuvant. This preparation is injected subcutaneously and is followed by booster injections of SP22 mixed with incomplete adjuvant. Sera obtained after the final booster injection are checked for titer, affinity, and specificity.

[0119.] Specifically, for rabbits, 100 micrograms of SP22 protein as obtained above, or peptides which are analogs of SP22, were solubilized in 0.5 mL physiological saline and emulsified with an equal volume of Freund's adjuvant to prepare inoculum sites in the back. Two New Zealand White female rabbits, weight 2.5-3.5 kg, were bled via the marginal ear vein for pre-immune serum. Approximately 50 microliters of inoculum was injected into 20 sites within the shaved area. The rabbits were boosted in similar fashion four weeks later. Two weeks later, the rabbits were bled again via the marginal ear vein, and sera containing the polyclonal antibodies was obtained.

097544-0404
"000000" 4559260

[0120.] To produce antiserum to SP22, a detergent extract of cauda epididymal sperm was chromatographed by reverse-phase HPLC and fractions enriched in SP22 were run in analytical two dimensional gels. Coomassie-stained SP22 punches were subsequently subjected to electroelution and electroeluted material was desalted, concentrated, and assayed for protein. After verifying that the electroeluted material was SP22, 25 micrograms were mixed with Freund's complete adjuvant and injected subcutaneously into each of six mice. Four other mice received only adjuvant.

[0121.] After four weeks, each mouse was boosted with 12.5 micrograms of SP22 mixed with Freund's incomplete adjuvant. After another ten days, a final similar booster was given. The mice were euthanized the days after the final booster, and the serum was collected.

[0122.] B. Monoclonal Antibody Production

BALB/c mice were immunized initially via i.p. injections with 50 ug of full length recombinant SP22 and later boosted similarly with native SP22. Services were provided by BioCon, Inc. (Rockville, MD). Spleens were collected and cell suspensions were prepared by perfusion with DMEM. The BLAB/c spleen cells were fused with SP 2/-Ag 14 mouse myeloma cells by PEG and the resultant hybridomas grown in HAT selected tissue culture media plus 20% fetal calf serum. The surviving cells were allowed to grow to confluence. The spent culture medium was checked for antibody titer, specificity, and

09544-0400

[0123.] Specifically, the mice were immunized with SP22 adjuvant emulsion described above. Each mouse first received 0.2 mL of this emulsion intraperitoneally, and then was reinfected in similar fashion with 0.1 mL six weeks later. Mouse serum was obtained then days after the second injection and then tested for anti-FRP activity via ELISA. The mouse exhibiting the highest possible anti- FRP activity was chosen for cell fusion.

[0125.] The next day, DMEM-FCS containing hypoxanthine, thymidine, and aminopterin (HAT medium) was added to each well. The concentration of HAT in the medium to be added was twice the final concentration required, i.e.,

$$H_{final} = 1 \times 10^{-4}M$$

$$A_{\text{final}} = 4 \times 10^{-7}\text{M}, \text{ and}$$

$$T_{\text{final}} = 1.6 \times 10^{-5}\text{M}.$$

[0126.] Subsequently, the plates were incubated with HAT medium every three to four days for two weeks. Fused cells were thereafter grown in DMEM-FCS containing hypoxanthine and thymidine. As cell growth became $\frac{1}{2}$ to $\frac{3}{4}$ confluent on the bottom of the wells, supernatant tissue culture fluid was taken and tested for SP22 specific antibody by ELISA. Positive wells were cloned by limiting dilution over macrophage or thymocyte feeder plates, and cultured in DMEM-FCS. Cloned wells were tested and recloned three times before a statistically significant monoclonal antibody was obtained. Spent culture media from the chosen clone contained antibody which bonds SP22 in all dilutions tested.

[0127.] C. Antibody to SP22 Peptides

SP22 is identified by its biological functions and activities set forth herein, as well as by its size of approximately 22 kD and isoelectric point of 5.25. However, changes in form and the substitution of fragments or equivalents are contemplated as circumstances may suggest or render expedient, including variations in methods for physically characterizing the protein. For instance, it may be necessary to generate polyclonal antibodies to peptide fragments of SP22 if sufficient amounts of purified SP22 cannot be obtained relatively easily.

[0128.] In addition to the antibodies which are identical to

the naturally-occurring SP22 peptide antibody, the present invention embraces epitopes which are substantially homologous with such antibodies.

[0129.] The term "antibody" is meant to include polyclonal antibodies, monoclonal antibodies, chimeric antibodies, anti-idiotypic antibodies to antibodies that can be labeled in soluble or bound form, as well as active fractions thereof provided by any known technique, such as, but not limited to, enzymatic cleavage, peptide synthesis, and recombinant techniques.

[0130.] Polyclonal antibodies are heterogeneous populations of antibody molecules derived from the sera of animals immunized with an antigen. A monoclonal antibody contains a substantially homogeneous population of antibodies specific to antigens, which population consists substantially similar epitope binding areas.

[0131.] Chimeric antibodies are molecules in which different proteins are derived from different animal species, such as those having the variable region derived from a murine monoclonal antibody and a human immunoglobulin constant region. Chimeric antibodies are primarily used to reduce immunogenicity in application and to increase yields in production, for example, where murine monoclonal antibodies have high yields from hybridomas but higher immunogenicity in human, such that human murine chimeric monoclonal antibodies are used.

[0132.] Chimeric antibodies and methods for their production are known in the art [Cabilly et al., Proc. Natl. Acad. Sci. USA 81:6851-6855 (1994); Boulilanne et al., Nature 312:643-646 (1984); European Patent Application 125023; Neiberger et al., Nature 314:2680279 (1985); Taniguchi et al., European Patent Application 171496; Morrison et al., European Patent Application 173494; Neuberger et al., PCT Patent Application WO 8601533; Kudo et al., European Patent Application 184187; Sahagah et al., J. Immunol. 137:1066-1074; Robinson et al., PCT Patent Application WO 8702671; Liu et al., Proc. Natl. Acad. Sci. USA 84:3439-3443 (1987); Sun et al., Proc. Natl. Acad. Sci. USA 84:214-218 (1987); Better et al., Science 240:1041-1043 (1988); and Harlow and Lane, Antibodies, a Laboratory Manual]. Each of these references is hereby incorporated herein by reference in its entirety.

[0133.] An anti-idiotypic antibody is an antibody which recognizes unique determinants generally associated with the antigen-binding site of an antibody. An anti-idiotypic antibody can be prepared by immunizing an animal of the same species and genetic type (e.g., a mouse strain) as the source of the monoclonal antibody with the monoclonal antibody to which an anti-idiotypic antibody is being prepared. The immunized animal will recognize and respond to the idiotypic determinants of the immunizing antibody by producing an antibody to these idiotypic determinants, i.e., the anti-idiotypic activity. See, for example, U.S. Patent No.

4,669,880, the entire contents of which are hereby incorporated by reference.

[0134.] The anti-idiotypic antibody may also be used as an immunogen to produce an immune response in yet another animal, producing a so-called anti-anti-idiotypic antibody. The anti-anti-idiotypic antibody may be epitopically identical to the original monoclonal antibody which induces the anti-idiotypic antibody. Thus, by using antibodies to the idiotypic determinants of a monoclonal antibody, it is possible to identify other clones expressing antibodies of identical specificity.

[0135.] Accordingly, monoclonal antibodies generated against SP22, and related proteins of the present invention, may be used to induce anti-idiotypic antibodies in suitable animals, such as BALB/c mice. Spleen cells from such immunized mice are used to produce anti-idiotypic hybridomas secreting anti-idiotypic monoclonal antibodies. Further, the anti-idiotypic monoclonal antibodies can be coupled to a carrier such as keyhole limpet hemocyanin (KLH) and used to immunize additional BALB/c mice. Sera from these mice will contain anti-anti-idiotypic antibodies that have the binding properties of the original monoclonal antibodies specific for SP22 or epitopes thereof.

[0136.] The term "antibody" is also meant to include both intact molecules as well as active fractions thereof, such as, for example, those which are capable of binding antigen. Fab

and F(ab')₂ fragments lack the Fc fragment of intact antibody, clear more rapidly from the circulation, and may have less non-specific tissue binding than an intact antibody, cf. Wahl et al., J. Nucl. Med. 24:316-325, 1983.

[0137.] The term "antagonist" includes antibody, complementary peptides or fragments thereof, or other small molecules which inhibit the activity of the protein.

[0138.] The term "substantially homologous", when used in connection with amino acid sequences, refers to sequences which are substantially identical to or similar in sequence with each other, giving rise to a homology of conformation and thus to retention, to a useful degree, of one or more biological (including immunological) activities. The term is not intended to imply a common evolution of the sequences.

[0139.] Substantially homologous peptide epitopes may be identified by a variety of techniques. It is known in the art that one may synthesize all possible single substitutions mutants of a known peptide epitope, Geysen et al., Proc. Nat. Acad. Sci. (USA) 18:3998-4002, 1984. While the effects of different substitutions are not always additive, it is reasonable to expect that two favorable or neutral single substitutions at different residue positions in the epitope can safely be combined in most cases.

[0140.] One may also synthesize a family of related single or multiple substitution mutants, present the mixture to a cell line capable of presenting the desired epitopes, and expose

057554-0001
FBI/DOJ

the cells to suitable restricted antigens. If the cells are lysed, effective epitopes may be identified either by direct recovery from the cells or by a progressive process of testing subsets of the effective peptide mixtures. Methods for the preparation of degenerate peptides are described in Rutter, U.S. Patent No. 5,010,175; Haughter et al., Proc. Nat. Acad. Sci. (USA) 82:5131-5135 (1985); Geysen et al., op. cit.; WO86/06487; and WO86/00991.

[0141.] In devising a multiple mutagenesis strategy, a person of ordinary skill would of course give weight to the single substitution mutant data in determining both which residues to vary and which amino acids or classes of amino acids are suitable replacements.

[0142.] It is also possible to predict substantially homologous epitopes by taking into account studies of sequence variations in families or naturally occurring homologous proteins. Certain amino acid substitutions are more often tolerated than others, and these are often correlated with similarities in size, charge, etc., between the original amino acid and its replacement. Insertions or deletions of amino acids may also be made, as described above.

Clones

[0143.] Once an antibody to, or portions sequence of, SP22 is available, it will be necessary to screen cDNA libraries to identify the cDNA clones that produce SP22. SP22 may be produced by methods other than recovery from male animals. In

0975644-040304
T02070-4735260

particular, a cDNA probe is prepared against a partial sequence of SP22 and used to identify the SP22 genome in cells from any mammalian species. The identified genome is then inserted into a plasmid which is then employed to produce recombinant SP22 in proliferating bacteria or other hosts according to methods known in the art. This will be useful in the methodologies, e.g., immunocontraception, addition to sperm, described herein. However, not only will these utilities require large amounts of SP22, they will also require large quantities of the SP22 antibody. This is accomplished in batch hybridoma cell culture using proven methods. The same processes can be used to identify antagonists to SP22 and fragments and derivatives thereof.

Contraception

[0144.] Antibodies to SP22 can be used for contraception as well as for assaying fertility. A reversible contraceptive vaccine is provided by administering to an animal subject SP22 as described above in an amount effective to reduce the fertility of that subject via generation of antibodies to SP22. Partial reduction in fertility, i.e., effects which are reflected as a reduction in fertility in a population of subjects, are intended as within the scope of the present invention.

[0145.] Any animal which expresses sperm surface SP22 may be treated by the immunocontraceptive method of the present invention, including both birds and mammals. Exemplary

0975254-040304
F0E0T04T525260

mammals include mice, rabbits, dogs, cats, cows, pigs, sheep, horses, and humans. Mammalian subjects are preferred. The vaccine can be administered to either females or males by any suitable means, including by intramuscular injection. The antibody can be administered topically, as by vaginal foam or by any convenient topical method in an appropriate carrier, e.g., by nasal spray.

[0146.] The term "protection", as used herein, is intended to include prevention or suppression of production of fertile sperm.

[0147.] It will be understood that in medicine, it is not always possible to distinguish between preventing and suppressing, since the ultimate inductive event or events may be unknown, latent, or the patient is not ascertained until well after the occurrence of the event or events. The term "protection", as used herein, is meant to include prophylaxis.

[0148.] The form of administration may be systemic or topical. For example, administration of such a composition may be by various parenteral routes such as subcutaneous, intravenous, intradermal, intramuscular, intraperitoneal, intranasal, transdermal, vaginal or buccal routes.

Alternatively, or concurrently, administration may be by the oral route. Parenteral administration can be by bolus injection or by gradual release over time.

[0149.] It is understood that the suitable dose of a composition according to the present invention will depend

RECEIVED 4-15-60

upon the age, health and weight of the recipient. However, the most preferred dosage can be tailored to the individual subject, as is understood and determinable by one skilled in the art, without undue experimentation. This typically involves adjustment of a standard dose, e.g., reduction of the dose if the patient has a low body weight.

[0150.] Prior to use in humans, a drug is first evaluated for safety and efficacy in laboratory animals. In human clinical trials, one begins with a dose expected to be safe in humans, based on the preclinical data for the drug in question, and on customary doses for analogous drugs, if any. If this dose is effective, the dosage may be decreased to determine the minimum effective dose, if desired. If this dose is ineffective, it will be cautiously increased, with the patients monitored for signs of side effects. See, e.g., Berkow et al., eds. The Merck Manual, 15th edition, Merck and Co., Rahway, NJ, 1987; Goodman et al., eds., Goodman and Gilman's The Pharmacological Basis of Therapeutics, 8th edition, Pergamon Press, Inc., Elmsford, NY, 1990; Avery's Drug Treatment: Principles and Practice of Clinical Pharmacology and Therapeutics, 3rd edition, ADIS Press, Ltd., Williams and Wilkins, Baltimore, MD, 1987; Ebadi, Pharmacology, Little Brown and Co., Boston, MA, 1985; which references and references cited therein are entirely incorporated herein by reference.

[0151.] The appropriate dosage form depends on the

[0152.] In addition to the protein or antigen of the invention, a pharmaceutical vaccine composition may contain suitable pharmaceutically acceptable carriers, such as excipients, carriers, and/or auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. The amount of antigen administered depends upon factors such as route of administration, species, and the use of booster administration. In general, dose of about 0.1 to about 100 micrograms per kg of body weight may be used. The antigen to SP22 may be prepared at both human and veterinary vaccine formulations. Vaccine formulations of the present invention comprise the antigen in a pharmaceutically acceptable carrier. The antigen is included in the carrier in an amount which is effective to reduce the fertility of the subject being treated. Pharmaceutically acceptable carriers are preferably liquid, particularly aqueous carriers, such as sodium phosphate buffered saline. The vaccine formulations may be stored in a sterile glass container sealed with a

[0153.] Vaccine formulations of the present invention may optionally contain one or more adjuvants. Any suitable adjuvant can be used, such as aluminum hydroxide, aluminum phosphate, plant and animal oils, and the like, with the amount of adjuvant depending on the nature of the particular adjuvant employed. In addition, the vaccine formulations may also contain at least one stabilizer, such as carbohydrates such as sorbitol, mannitol, starch, sucrose, dextrin, and glucose, as well as proteins such as albumin or casein, and buffers such as alkali metal phosphates and the like.

Preferably, the preparations contain from about 0.1 to about 99 percent, preferably from about 25 to 85 percent, of active ingredient, together with the excipients. The excipients may be any pharmaceutically acceptable excipients or carrier which can be used with the antigen or antibody.

[0155.] Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, and sorbitol, cellulose preparations and derivatives and/or calcium phosphates. Also useful as excipients are binders such as

[0156.] For vaginal application, suppositories, lotions, creams, sprays, or foams may be used to incorporate the active ingredients. Suitable suppository bases are, for example, natural or synthetic triglycerides, paraffin hydrocarbons, polyethylene glycols or higher alkanols. Foam formulations may include oily suspensions or aqueous solutions of the active ingredient with suitable foaming agents. Other topical carriers for vaginal applications include pharmaceutically acceptable liquids in which the active ingredient is suspended or dissolved.

[0158.] SP22 can also be used to identify male animals who are good candidates for supplying sperm for artificial insemination. Since many livestock animals are reproduced by artificial insemination or embryo transfer, it is important to be able to identify males who are fertile as well as possessing desirable characteristics to pass on to the next generation. Techniques for reproducing animals by embryo transfer are described in U.S. Patents 3,854,479; 4,816,257;

and 4,326,505, the entire contents of each of which are hereby incorporated by reference. By determining the amount of SP22 in the sperm of a subject animal, the fertility of the animal can be predicted.

[0159.] It is known that sperm proteins are affected by toxicants and pollutants. According to the present invention, the changes in SP22 level are calibrated to predict the likelihood of the sperm having been rendered infertile because of exposure to the toxicant.

[0160.] In Klinefelter et al., 1996, it was demonstrated that endocrine-disruptive chemicals decreased the fertilizing ability of cauda epididymal sperm in four days. Tests were conducted to determine if this infertility was related to decrease of SP22 associated with the sperm.

[0161.] In order to evaluate the effects of exposure to toxicants that perturb the androgen status of the animal, such as EDS, sperm are subjected to analysis of SP22. Adult (90 to 120 day old) male Sprague-Dawley rats were housed two to three per cage with laboratory-grade pine shavings as bedding. The rats were maintained under controlled temperature (22°C) and humidity (40-50%) conditions, and were given Purina laboratory rat chow and tap water *ad libitum*. Males were maintained in a 14-hour light, 10-hour dark schedule. Each male was numbered and randomly assigned to a treatment group. The test toxicant was administered either as a single intraperitoneal injection or as four daily injections. After four days the rats were

09:34:04.0404

100-4523260
killed, and the caudal epididymides of each rat was placed in a 35-mm culture dish containing 2 mL of Medium 199. Detergent extracts representing $10\text{--}40 \times 10^6$ sperm, depending on treatment, were prepared and aliquots equivalent to 30 micrograms were electrophoresed in a mini, two dimensional electrophoresis system (BioRad) for quantitative analysis of SP22. Specifically, sperm were transferred to a microcentrifuge tube and washed twice by centrifugation ($300 \times g$, five minutes) in Dulbecco's phosphate buffered saline, pH 7.2, with freshly added 0.2 extracted phenylmethylsulfonyl fluoride (PMSF). After the final wash, the sperm were extracted for one hour at room temperature with 1 mL of 40 extracted n-octyl- β - glucopyranoside in 10 ml extracted Tris, pH 7.2, containing freshly added PMSF. Following a final centrifugation at $3000 \times g$, the supernatant was removed and frozen at -70°C .

[0162.] Upon thawing, each extract was concentrated in 1 extracted Tris buffer by two centrifugations ($3000 \times g$ for 45 minutes at 4°C) in Centricon-10 units (Amincon). Protein concentration was determined using a Pierce protein assay kit. Sample volumes containing 30 μg protein were lyophilized, and protein was solubilized for 30 minutes at room temperature in 45 μL of sample buffer consisting of 5.7 g urea, 4 mL 10% NP-40, 0.5 mL ampholytes (70% 3-10, 30% 5-7) and 0.1 g dithiothreitol per 10 mL. Isoelectric focusing (750 V, 3.5 hours) was conducted in gels consisting of 6.24 g. urea, 1.5 g

acrylamide (30% acrylamide, 1.2% bisacrylamide), 2.25 mL 10% NO-40, and 0.65 mL ampholytes (60% 3-10, 40% 5-7) per 10 mL. Molecular weight separation was conducted in 11% methanol and silver stained. A Kepler two dimensional gel analysis system (Large Scale Biology Corp., Rockville, MD) was used for background correction, spot matching, and spot area quantitation. Images were acquired by transmitters at 80 μ m spatial resolution and 4096 gray levels on an Ektron 1412 scanner and converted to 256 gray levels. Quantitation was done by fitting two-dimensional Gaussian distributions to the density distribution of the spot area following background subtraction. Of the 124 proteins (spots) that were identified in the 50 gel data set, 22 were common to gels representative of sperm extracts of vehicle-treated animals. Of these 22 proteins, only SP22 was affected by all test chemicals in a dose-related fashion. In fact, SP22 was the only one of the 124 that were identified, that changed in either a dose or treatment-related fashion.

[0163.] It has been determined that insemination (*in utero*) of 5×10^6 epididymal sperm from a control rat results in approximately 75% fertility, thereby providing relatively greater sensitivity than insemination of a number of sperm that would result in 100% fertility.

[0164.] The various data (fertility and SP22, as well as other endpoints such as motility parameters and testosterone concentrations) were collected and analyzed using two-way

analysis of variance for both block and treatment effects. An initial analysis was performed to determine whether experimental block differences influenced the parameters measured. Where overall block effects are significant ($p < 0.05$), the least-square means were compared for significant ($p < 0.05$) treatment differences. A correlation analysis was performed to determine whether significant ($p < 0.01$) correlations exist between each of the measured endpoints, and fertilizing ability and Pearson correlation coefficients (R) were calculated.

[0165.] In a study reported in Klinfelter et al., Journal of Andrology 14(4):318-327, 1994, the authors used *in utero* insemination of epididymal sperm and exposure to a chemical which disrupts androgen status of the epididymis, ethane dimethanesulfonate, EDS, to investigate the hypothesis that EDS compromises the fertilizing ability of sperm by affecting epididymal function directly. Fertilizing ability, sperm motility, serum testosterone, and tissue testosterone were evaluated. In addition, sperm proteins were extracted and analyzed by quantitative two dimensional gel electrophoresis. An 18 kD protein was well correlated with fertility. However, it was felt that changes in this protein were not sufficient either to EDS itself or the dose that was tested.

[0166.] In a subsequent study, the insemination procedure was modified to permit assessment of fertility (implants/corpora lutea) rather than fertilizing ability

(percentage of eggs fertilized). In this study, multiple chemicals that disrupt endocrine status were tested. Adult males were exposed either to 25 or 50 mg/kg EDS, epichlorohydrin, 3 or 6 mg/kg, or hydroflutamide, 12.5 or 25.0 mg/kg, or chloroethylmethanesulfonate, 12.5 or 18.75 mg/kg. Each of these compounds perturbs the endocrine balance of the male reproductive system. The animals exposed to the known antiandrogen hydroglutamate were castrated and implanted with testosterone implants just prior to the first injection. The vehicle controls for all treatments except hydroxyflutamide received daily injections of 30% DMSO in water. The vehicle controls for the hydroxyflutamide animals were castrated, implanted with testosterone implants, and given daily injection of 15% ethanol. Four days after the onset of dosing, the males were killed and the epididymides were removed. The caput-corpus was frozen on dry ice for subsequent steroid extraction and testosterone assay. Sperm were released from the epididymal tubule into insemination medium and held in CO₂ incubator at 34°C for no more than 15 minutes until insemination. Adult, estrus-synchronized female rats were monitored for lordosis behavior just after lights out on the day of insemination. Females displaying mating behavior were cervically stimulated with vasectomized teaser males at least 15 minutes prior to insemination. A volume equal to 5×10^6 sperm was inseminated into each uterine horn at day 0. On day 9, the females were killed and fertility was

assessed. A Kepler two dimensional gel analysis system (Large Scale Biology Corp., Rockville, MD) was used for background correlations, spot matching, and spot area quantitation. Images were acquired by transmittance at 80 μ m spatial resolution and 4096 gray levels. Quantitation was done by fitting two-dimensional Gaussian distributions to the density distribution of the spot area following background subtraction. Of the 125 proteins (spots) that were identified in the 50 gel data set, 22 were common to gels representative of sperm extracts of vehicle-treated animals. Of these 22 proteins, only SP22 was affected by all test chemicals in a dose-related fashion. In fact, SP22 was the only one, of the 124 that were identified, that changed in either a dose or treatment-related fashion.

[0167.] Measurements of sperm motion and sperm morphology were not significantly affected by any of the treatments. Based on scatter plot of the data relating the amount of SP22 to fertility (frequent), fertility classes greater and less than n 50% were chosen. Variables were then entered into the discriminant analyst to predict fertility by class, as shown in Table 1. Since, in this study, fertility for the control animals was targeted at $68\% \pm$ a standard deviation of 18%; 50% represented a reasonable cutoff for the fertile class.

Table 1
Discrimination Analysis Based on SP22

CLASS	PERCENTAGE CORRECTLY PREDICTED
Fertile (>50%)	90 (17/19)
Subfertile (<50%)	94 (29/31)

[0168.] A regression analysis showed that the amount of SP22 was significantly correlated to fertility ($p < 0.0001$; $r^2 = 0.83$). A nonlinear fit of the data was indicated, since a threshold of 10,000 integrated optical density units of SP22 was necessary to achieve greater than 50% fertility.

[0169.] Thus, by entering the level of SP22 of a sperm sample into an appropriate mathematical model, it is possible to predict the fertility of the sperm sample with a reasonably high degree (i.e., >90%) of success. An antibody to SP22 can be used to evaluate the fertility of sperm in an epididymal sperm sample or an ejaculate. Since the antibody to SP22 recognizes a single protein on immunoblots of cells of both human and stallion sperm extracts, this antibody will most likely be applicable to evaluation of animals in which maximum fertility is important, e.g., cattle, horses, dogs, and humans among other animals.

[0170.] The toxicants tested above do perturb the endocrine balance of the male reproductive system. Other environmentally relevant endocrine disruptors, such as dioxin, could also compromise the expression of SP22. The present invention thus includes a screening kit to test such chemicals.

[0171.] The foregoing description of the specific embodiments will so fully reveal the general nature of the invention that others can, by applying current knowledge, readily modify and/or adapt for various application such specific embodiments without departing from the generic concept, and therefore such adaptations and modifications are intended to be comprehended within the meaning and range of equivalents of the disclosed embodiments. It is to be understood that the phraseology or terminology herein is for the purpose of description and not of limitation.

[0172.] All references cited in this specification are hereby incorporated by reference.

097554 010301

REFERENCES

Amann, et al. Responses of Mammalian Sperm Exposed to Synthetic FertPlus™ Peptide. J. Androl. Suppl.: 102A (1998a).

Amann, et al. Exposure of Human, Bull, or Boar Sperm to a Synthetic Peptide Increases Binding to an Egg-Membrane Substrate. J. Andro. (Submitted, 1998b).

Benton, et al. Screening λ gt Recombinant Clones by Hybridization to Single Plaques *in situ*. Science, 196:180 (1977).

Burks, et al. Interaction of a Tyrosine Kinase from Human Sperm with the Zona Pellucida at Fertilization. Science, 269:83-89 (1995).

Cohen, et al. Mammalian Sperm-Egg Fusion: The Development of Rat Oolemma Fusibility during Oogenesis Involves the Appearance of Binding Sites for Sperm Protein "DE". Biol. Reprod. 55:200-206 (1996).

Capkova, et al. Binding of Boar Spermatozoa to Porcine Oocytes: Effect of Low Molecular Weight 170kDa Protein. Mol. Reprod. Devel. 46:168-175 (1997).

Causnicu, et al. Antibodies Against Epididymal Glycoproteins Block Fertilizing Ability in Rat. J. Reprod. Fert. 72:467-471 (1984).

Gerena, et al. Identification of a Fertility-Associated Protein in Bull Seminal Plasma as Lipocalin-Type Prostaglandin D Synthase. Biol. Reprod. 58:826-833 (1998).

Hammerstedt, et al. A Method and Use of Polypeptide in Sperm-Egg Binding to Enhance or Decrease Fertility. International Patent Publication #WO/97/25620 (1997).

Hunnicut, et al. Sperm Surface Protein PH-20 is Bifunctional: One Activity is a Hyaluronidase and a Second, Distinct Activity is Required in Secondary Sperm-Zona Binding. Biol. Reprod. 55:80-86 (1996).

Killian, et al. Fertility-Associated Proteins in Holstein Bull Seminal Plasma. Biol. Reprod. 49:1202-1207 (1993).

Killian, et al. Alteration and Prediction of Male Fertility using Seminal Plasma and its Components. United States Patent #5,569,581 (1996).

11/11/97 11:11:11

Klinefelter, et al. Discriminant Analysis Indicates a Single Sperm Protein (SP22) is Predictive of Fertility Following Toxicant Exposure. J. Andrology, 18:139-150 (1997).

Klinefelter, et al. Reduced Fertility is Correlated with SP22 Levels Following Exposure to the Disinfection Byproduct of Drinking Water, Bromochloroacetic Acid. J. Androl. (Manuscript in Prep.)

Lea, et al. Cloning and Sequencing of cDNAs Encoding the Human Sperm Protein, Sp17. Biochimica. Biophysica Acta, 1307:263-266 (1996).

Linder, et al. Dibromoacetic Acid Affects Reproductive Competence and Sperm Quality in the Male Rat. Fund. Appl. Toxicol. 28:9-17 (1995).

Linder, et al. Histopathologic Changes in the Tests of Rats Exposed to Dibromoacetic Acid. Reprod. Toxicol. 11:47-56 (1997).

Linder, et al. Spermatotoxicity of Dichloroacetic Acid. Reprod. Toxicol. 11:681-688 (1997).

Wei., et al. Fertility Studies with Antisperm Antibodies. Arch. Androl. 32:251-262 (1994).

2025 RELEASE UNDER E.O. 14176

Nagakubo, et al. A Novel Oncogene Which Transforms Mouse NIH3T3 Cells in Cooperation with Ras. Biochem. Biophys. Res. Comm. 231:509-513 (1997).

O'Rand, et al. Monoclonal Antibodies to Rabbit Sperm Autoantigens. I. Inhibition of In Vitro Fertilization and Localization on the Egg. Biol. Reprod. 30:721-730 (1984).

O'Rand, et al. Sperm Antigen Corresponding to a Sperm Zona Binding Protein Autoantigenic Epitope. United States Patent #5,480,799 (1996).

Primakoff, et al. Reversible Contraceptive Effect of PH-20 Immunization in Male Guinea Pigs. Biol. Reprod. 56:1142-1146 (1997).

Peknicova, et al. Binding of Boar Spermatozoa to Porcine Oocytes: Effect of Low Molecular Weight 17-kDa Protein. Mol. Reprod. Devel. 46:168-175 (1997).

Sanger, et al. DNA Sequencing with Chain Terminating Inhibitors. Proc. Natl. Acad. Sci. USA 74:5463-5467 (1977).

Welch, et al. Expression of a Glyceraldehyde 3-phosphate Dehydrogenase Gene Specific to Mouse Spermatogenic Cells. Biol. Reprod. 46:869-878 (1992).

2025-04-15 15:00

Welch, J.E Barbee, R.R., Roberts, N.L, Suarez, J.D., and Klinefelter, G.R., SP22: a Novel Fertility Protein from a Highly Conserved Gene Family. J. Andrology 19:395-398 (1998)

Klinefelter, G.R. and Welch, J.E., The Saga of a Male Fertility Protein (SP22). Annual Review of Biomedical Sciences 1:145-184 (1999).

Eidne, K.A., Henery, C.C., and Aitken, R.J. Selection of Peptides Targeting the Human Sperm Surface using Random Peptide Phage Display to Identify Ligands Homologous to ZP. Biol. Reprod. 63:1396-1402 (2000).

O'Rand, MG and Widgren, EE, Identification of Sperm Antigen Targets for Immunocontraception: B-Cell Epitope Analysis for SP17. Reprod. Fertil. Dev. 6: 189-196 (1994).

2025 RELEASE UNDER E.O. 14176